

# The dark halo of Milky Way-like galaxies

Nassim Bozorgnia




Institute for Particle Physics Phenomenology  
Durham University

# Evidence for Dark Matter

## Galaxy rotation curves

## Dwarf galaxies




## Galaxy clusters



## Gravitational lensing


## Cosmic Microwave Background

## Large Scale Structure




# Cosmic Microwave Background

Measurements of temperature fluctuations in the CMB provide a precise determination of the Dark Matter (DM) density in the Universe.




Planck 2015



# Our simulated Universe

# Dark Matter halo


*What is the distribution of DM in halo of our Galaxy?*



# Dark Matter halo



*What is the distribution of DM in halo of our Galaxy?*

Uncertainties in the DM distribution → *prevents a precise determination of the properties of the DM particle.*




# Dark Matter searches

- **WIMPs** are the most extensively studied class of DM candidates, and can be searched for in three complimentary ways:




# Dark Matter searches

- **WIMPs** are the most extensively studied class of DM candidates, and can be searched for in three complimentary ways:





- 
- Use high resolution cosmological simulations to extract the DM distribution and make accurate predictions for **direct** and **indirect** DM searches.


# Prospects for direct DM searches

# Dark Matter direct detection

- Search for WIMPs by measuring the recoil energy of a nucleus in an underground detector after collision with a WIMP.

- Elastic recoil energy:

$$E_R = \frac{2\mu_{\chi N}^2 v^2}{m_N} \cos^2 \theta$$



- Minimum WIMP speed required to produce a recoil energy  $E_R$  :

$$v_{\min} = \sqrt{\frac{m_N E_R}{2\mu_{\chi N}^2}}$$

# Direct detection event rate

- The differential event rate (per unit detector mass):

$$\frac{dR}{dE_R} = \frac{\rho_\chi}{m_\chi m_N} \int_{v > v_{\min}} d^3v \frac{d\sigma_{\chi N}}{dE_R} v f_{\text{det}}(\mathbf{v}, t)$$

# Direct detection event rate

- The differential event rate (per unit detector mass):

$$\frac{dR}{dE_R} = \frac{\rho_\chi}{m_\chi m_N} \int_{v > v_{\min}} d^3v \frac{d\sigma_{\chi N}}{dE_R} v f_{\text{det}}(\mathbf{v}, t)$$

astrophysics

- Astrophysical inputs:**
  - local DM density:** *normalization in event rate.*
  - local DM velocity distribution:** *enters the event rate through an integration.*

# Direct detection event rate

- The differential event rate (per unit detector mass):

$$\frac{dR}{dE_R} = \frac{\rho_\chi}{m_\chi m_N} \int_{v > v_{\min}} d^3v \frac{d\sigma_{\chi N}}{dE_R} v f_{\text{det}}(\mathbf{v}, t)$$

astrophysics

- For standard spin-independent and spin-dependent interactions:

$$\frac{d\sigma_{\chi N}}{dE_R} = \frac{m_N}{2\mu_{\chi N}^2 v^2} \sigma_0 F^2(E_R)$$

# Direct detection event rate

- The differential event rate (per unit detector mass):

$$\frac{dR}{dE_R} = \frac{\rho_\chi}{m_\chi m_N} \int_{v > v_{\min}} d^3v \frac{d\sigma_{\chi N}}{dE_R} v f_{\text{det}}(\mathbf{v}, t)$$

astrophysics

- For standard spin-independent and spin-dependent interactions:

$$\frac{dR}{dE_R} = \frac{\sigma_0 F^2(E_R)}{2m_\chi \mu_{\chi N}^2} \rho_\chi \eta(v_{\min}, t)$$

particle physics



astrophysics

where

$$\eta(v_{\min}, t) \equiv \int_{v > v_{\min}} d^3v \frac{f_{\text{det}}(\mathbf{v}, t)}{v}$$

Halo integral

# Direct detection status



- Assumption in these kinds of plots: **Standard Halo Model (SHM)**



# Standard Halo Model


- The simplest model for the DM distribution in our Galaxy is the *Standard Halo model*: isothermal sphere with an isotropic Maxwell-Boltzmann velocity distribution.

Drukier, Freese, Spergel, 1986

# Standard Halo Model

- The simplest model for the DM distribution in our Galaxy is the *Standard Halo model*: isothermal sphere with an isotropic Maxwell-Boltzmann velocity distribution.


Drukier, Freese, Spergel, 1986

- Hydrostatic equilibrium: pressure balances gravitational potential
  - Density profile:  $\rho(r) \propto r^{-2}$
  - Local DM density:  $0.3 \text{ GeV/cm}^3$
  - Typical DM speed: 220 km/s
- 
- Actual DM distribution may *deviate substantially* from the SHM.

# Local Dark Matter density

## From observations:



- **Local estimates:** use kinematical data from a nearby population of stars.
- Robust measurements, but need to account for the local contribution of baryons which has significant uncertainties. → *large error bars*




# Local Dark Matter density

## From observations:

- **Local estimates:** use kinematical data from a nearby population of stars.
  - Robust measurements, but need to account for the local contribution of baryons which has significant uncertainties. → *large error bars*
- **Global estimates:** based on mass modeling of the MW, and fits to kinematical data across the Galaxy.
  - Good precision ( $\sim 10\%$ ), but estimates are strongly model dependent. → *systematic uncertainties*



# Local Dark Matter density



Read, 1404.1938

# Local DM velocity distribution

- The velocity distribution depends on the halo model.
- In the **SHM**, a truncated Maxwellian velocity distribution is assumed:

$$f_{\text{gal}}(\mathbf{v}) = \begin{cases} N \exp(-\mathbf{v}^2/v_c^2) & v < v_{\text{esc}} \\ 0 & v \geq v_{\text{esc}} \end{cases}$$

with  $v_c = 220$  km/s and  $v_{\text{esc}} = 550$  km/s.

$\sigma_v = \sqrt{3/2} v_c$  independent of radius.

# Local DM velocity distribution

- The velocity distribution depends on the halo model.
- In the **SHM**, a truncated Maxwellian velocity distribution is assumed:

$$f_{\text{gal}}(\mathbf{v}) = \begin{cases} N \exp(-\mathbf{v}^2/v_c^2) & v < v_{\text{esc}} \\ 0 & v \geq v_{\text{esc}} \end{cases}$$


with  $v_c = 220$  km/s and  $v_{\text{esc}} = 550$  km/s.

$\sigma_v = \sqrt{3/2} v_c$  independent of radius.

- What can we learn from numerical simulations of galaxy formation about the local DM velocity distribution?

# Dark Matter only simulations

- DM speed distributions from cosmological N-body simulations **without baryons**, deviate substantially from a Maxwellian.



$$f(|\mathbf{v}|) = v^2 \int d\Omega_{\mathbf{v}} f(\mathbf{v})$$


Vogelsberger et al., 0812.0362

- Significant systematic uncertainty since the impact of baryons neglected.*



# Hydrodynamical simulations

- Each hydrodynamical (**DM + baryons**) simulation adopts a different *galaxy formation model, spatial resolution, DM particle mass*.




Bozorgnia & Bertone, 1705.05853


- Large variation in DM speed distributions between the results of different simulations.

# Hydrodynamical simulations

- Each hydrodynamical (**DM + baryons**) simulation adopts a different *galaxy formation model, spatial resolution, DM particle mass*.



Different criteria used to identify MW-like galaxies among different groups. The most common criteria is the MW mass constraint, which has a large uncertainty.



Bozorgnia & Bertone, 1705.05853

- Large variation in DM speed distributions between the results of different simulations.

# Hydrodynamical simulations

- To make precise quantitative predictions:
  - Model baryonic processes in a way that the main galaxy population properties are broadly reproduced.
  - Identify MW-like galaxies by taking into account *observational constraints on the MW*.

# Hydrodynamical simulations

- We use the **EAGLE** and **APOSTLE** hydrodynamic simulations.

Name	L (Mpc)	N	$m_g (M_{\text{sun}})$	$m_{\text{DM}} (M_{\text{sun}})$
<b>EAGLE HR</b>	25	$8.5 \times 10^8$	$2.26 \times 10^5$	$1.21 \times 10^6$
<b>APOSTLE IR</b>	—	—	$1.3 \times 10^5$	$5.9 \times 10^5$

- APOSTLE IR**: zoomed simulations of Local Group-analogue systems, comparable in resolution to **EAGLE HR**.


# Hydrodynamical simulations

- We use the **EAGLE** and **APOSTLE** hydrodynamic simulations.

Name	L (Mpc)	N	$m_g (M_{\text{sun}})$	$m_{\text{DM}} (M_{\text{sun}})$
<b>EAGLE HR</b>	25	$8.5 \times 10^8$	$2.26 \times 10^5$	$1.21 \times 10^6$
<b>APOSTLE IR</b>	—	—	$1.3 \times 10^5$	$5.9 \times 10^5$



- APOSTLE IR**: zoomed simulations of Local Group-analogue systems, comparable in resolution to **EAGLE HR**.
- Calibrated to reproduce the observed distribution of stellar masses and sizes of low-redshift galaxies.*
- Companion Dark Matter only (DMO) simulations were run assuming all the matter content is collisionless.

# EAGLE Simulations




**EAGLE Simulations, I407.7040**

# Milky Way analogues



# Identifying Milky Way analogues

- Identify MW-like galaxies by taking into account observational constraints on the MW, in addition to the mass constraint:  
**rotation curves** [Iocco, Pato, Bertone, 1502.03821], **total stellar mass**.




Bozorgnia et al., 1601.04707  
Calore, Bozorgnia et al., 1509.02164



# Identifying Milky Way analogues


- Identify MW-like galaxies by taking into account observational constraints on the MW, in addition to the mass constraint:  
**rotation curves** [Iocco, Pato, Bertone, 1502.03821], **total stellar mass**.



Bozorgnia et al., 1601.04707  
Calore, Bozorgnia et al., 1509.02164


# Dark Matter density profiles

- Spherically averaged DM density profiles of the MW analogues:



# Dark Matter density profiles


- Spherically averaged DM density profiles of the MW analogues:



- To find the DM density at the position of the Sun, consider a torus aligned with the stellar disc.



$$\rho_\chi = 0.41 - 0.73 \text{ GeV/cm}^3$$

Bozorgnia et al., 1601.04707



# Local Dark Matter density

Is there an enhancement of the local DM density in the Galactic disk compared to the halo?




- $\rho_{\text{torus}}$  larger than  $\rho_{\text{shell}}$  by 2 – 27% for 10 haloes.
- The increase in the DM density in the disk could be due to the DM halo contraction as a result of dissipational baryonic processes.

# Halo shapes

- To study the shape of the inner ( $R < 8$  kpc) DM haloes, calculate the inertia tensor of DM particles within 5 and 8 kpc.  
→ ellipsoid with three axes of length:


$$a \geq b \geq c$$



# Halo shapes

- To study the shape of the inner ( $R < 8$  kpc) DM haloes, calculate the inertia tensor of DM particles within 5 and 8 kpc.  
→ ellipsoid with three axes of length:

$$a \geq b \geq c$$




- **Sphericity:**  $s = c/a$  ( $s = 1$  : perfect sphere)
- **Hydro haloes:** at 5 kpc,  $s = [0.85, 0.95]$ . At 8 kpc,  $s$  lower by less than 10%.
- **DMO haloes:**  $s = [0.75, 0.85]$
- Due to dissipational baryonic processes, DM sphericity systematically higher in the hydro compared to DMO haloes.

# Halo shapes


- Describe a deviation from a sphere by the triaxiality parameter:

$$T = \frac{a^2 - b^2}{a^2 - c^2}$$

- Oblate systems:  $a \approx b \gg c \rightarrow T \approx 0$



- Prolate systems:  $a \gg b \approx c \rightarrow T \approx 1$




# Halo shapes


- Describe a deviation from a sphere by the triaxiality parameter:

$$T = \frac{a^2 - b^2}{a^2 - c^2}$$

- Oblate systems:  $a \approx b \gg c \rightarrow T \approx 0$



- Prolate systems:  $a \gg b \approx c \rightarrow T \approx 1$





- In the hydro case, inner haloes very close to spherical and deviation towards either oblate or prolate is small. **DMO counterparts** have a preference for *prolate inner haloes*.



# Local speed distributions


In the galactic rest frame:



Bozorgnia et al., 1601.04707

# Local speed distributions

In the galactic rest frame:



Bozorgnia et al., 1601.04707

- Maxwellian distribution with a free peak provides a better fit to haloes in the hydrodynamical simulations compared to their DMO counterparts.
- Best fit peak speed:  $v_{\text{peak}} = 223 - 289 \text{ km/s}$


# Local speed distributions

## Common trends in different hydrodynamical simulations:

- Baryons deepen the gravitational potential in the inner halo, shifting the peak of the DM speed distribution to *higher speeds*.
- In most cases, baryons appear to make the local DM speed distribution *more Maxwellian*.

Bozorgnia & Bertone, 1705.05853


# Departure from isothermal



Bozorgnia & Bertone, 1705.05853


- At the Solar circle, haloes in the hydrodynamical simulation are closer to isothermal than their DMO counterparts.

# Components of the velocity distribution



Bozorgnia et al., I601.04707

# Comparison with DMO



Bozorgnia et al., 1601.04707

# How common are dark disks?

- Clear velocity anisotropy at the Solar circle.
- Two haloes have a rotating DM component in the disc with mean velocity comparable (within 50 km/s) to that of the stars.

# How common are dark disks?

- Clear velocity anisotropy at the Solar circle.
- Two haloes have a rotating DM component in the disc with mean velocity comparable (within 50 km/s) to that of the stars.
- Hint for the existence of a co-rotating dark disk in 2 out of 14 MW-like haloes. → Dark disks are relatively rare in our halo sample.

Bozorgnia et al., 1601.04707

Schaller et al., 1605.02770



# How common are dark disks?

- Clear velocity anisotropy at the Solar circle.
- Two haloes have a rotating DM component in the disc with mean velocity comparable (within 50 km/s) to that of the stars.
- Hint for the existence of a co-rotating dark disk in 2 out of 14 MW-like haloes. → Dark disks are relatively rare in our halo sample.


Bozorgnia et al., 1601.04707

Schaller et al., 1605.02770

- *Sizable dark disks also rare in other hydro simulations:*
  - They only appear in simulations where a large satellite merged with the MW in the recent past, which is robustly excluded from MW kinematical data.

Bozorgnia & Bertone, 1705.05853


# The halo integral



- Halo integrals for the best fit Maxwellian velocity distribution (*peak speed 223 - 289 km/s*) fall within the  $1\sigma$  uncertainty band of the halo integrals of the simulated haloes.

Bozorgnia et al., 1601.04707

# The halo integral



- Baryons affect the velocity distribution strongly at the Solar position, resulting in a shift of the tails of the halo integrals to higher velocities with respect to DMO.

Bozorgnia et al., 1601.04707

# The halo integral

Common trend in different hydrodynamical simulations:

- Halo integrals and hence direct detection event rates obtained from a **Maxwellian velocity distribution with a free peak** are similar to those obtained directly from the simulated haloes.

Bozorgnia et al., [1601.04707](#) (EAGLE & APOSTLE)


Kelso et al., [1601.04725](#) (MaGICC)

Sloane et al., [1601.05402](#)

Bozorgnia & Bertone, [1705.05853](#)


# Implications for direct detection

- Assuming the **Standard Halo Model**:




# Implications for direct detection

- Compare with simulated Milky Way-like haloes:




# Implications for direct detection



- Difference in the local DM density  $\rightarrow$  overall difference with the SHM.
- Variation in the peak of the DM speed distribution  $\rightarrow$  shift in the low mass region.

# Implications for direct detection

Comparison to other hydrodynamical simulations:



Bozorgnia & Bertone, 1705.05853



# Non-standard interactions

- For a very general set of non-relativistic effective operators:

Kahlhoefer & Wild, 1607.04418

$$\frac{d\sigma_{\chi N}}{dE_R} = \frac{d\sigma_1}{dE_R} \frac{1}{v^2} + \frac{d\sigma_2}{dE_R}$$

# Non-standard interactions

- For a very general set of non-relativistic effective operators:

Kahlhoefer & Wild, 1607.04418

$$\frac{d\sigma_{\chi N}}{dE_R} = \frac{d\sigma_1}{dE_R} \frac{1}{v^2} + \frac{d\sigma_2}{dE_R}$$

$\eta(v_{\min}, t)$        $h(v_{\min}, t) = \int_{v > v_{\min}} d^3v \, v \, f_{\text{det}}(\mathbf{v}, t)$


# Non-standard interactions

- For a very general set of non-relativistic effective operators:

Kahlhoefer & Wild, 1607.04418

$$\frac{d\sigma_{\chi N}}{dE_R} = \frac{d\sigma_1}{dE_R} \frac{1}{v^2} + \frac{d\sigma_2}{dE_R}$$

$\eta(v_{\min}, t)$        $h(v_{\min}, t) = \int_{v > v_{\min}} d^3v v f_{\text{det}}(\mathbf{v}, t)$



Bozorgnia & Bertone, 1705.05853

- Best fit Maxwellian  $h(v_{\min})$  falls within the  $1\sigma$  uncertainty band of the  $h(v_{\min})$  of the simulated haloes.

# Dark Matter substructure


- High resolution **DMO** simulations predict:
  - DM density at the Solar position very smooth. Chance of the Sun residing in a DM subhalo of any mass is  $10^{-4}$ .  
[Vogelsberger et al., 0812.0362](#)
  - DM streams at the Solar position are unlikely to be important.  
[Vogelsberger & White, 1002.3162](#)

# Dark Matter substructure

- High resolution **DMO** simulations predict:
  - DM density at the Solar position very smooth. Chance of the Sun residing in a DM subhalo of any mass is  $10^{-4}$ .  
**Vogelsberger et al., 0812.0362**
  - DM streams at the Solar position are unlikely to be important.  
**Vogelsberger & White, 1002.3162**
- **What happens when baryons are included?**  
Substructure abundance reduced. **Sawala et al., 1609.01718**  
**Garrison-Kimmel et al., 1701.03792**  
*Need higher resolution hydro simulations to probe Solar position.*

# Dark Matter substructure


- High resolution **DMO** simulations predict:
  - DM density at the Solar position very smooth. Chance of the Sun residing in a DM subhalo of any mass is  $10^{-4}$ .  
*Vogelsberger et al., 0812.0362*
  - DM streams at the Solar position are unlikely to be important.  
*Vogelsberger & White, 1002.3162*
- **What happens when baryons are included?**  
Substructure abundance reduced. *Sawala et al., 1609.01718*  
*Garrison-Kimmel et al., 1701.03792*  
*Need higher resolution hydro simulations to probe Solar position.*
- **Promise of Gaia:** detect many stellar streams in the Solar neighborhood. Interaction of **DM subhaloes** and **stellar streams** can cause perturbations in the streams.  
*Detectable in future Gaia data!*  
*N. Banik, G. Bertone, J. Bovy and N. Bozorgnia, in preparation*



# Prospects for indirect DM searches

# Dark Matter indirect detection

- Search for Standard Model particles produced by the annihilation or decay of DM.






# Dark Matter indirect detection

- Expected gamma-ray flux from DM annihilation:

$$\frac{d\Phi_\gamma}{dE} = \frac{\langle\sigma v\rangle}{8\pi m_\chi^2} \frac{dN_\gamma}{dE} \int_{\text{l.o.s.}} ds \rho^2(r(s, \psi))$$



Pieri et al, 0908.0195


# Dark Matter indirect detection

- Expected gamma-ray flux from DM annihilation:

$$\frac{d\Phi_\gamma}{dE} = \frac{\langle\sigma v\rangle}{8\pi m_\chi^2} \frac{dN_\gamma}{dE} \int_{\text{l.o.s.}} ds \rho^2(r(s, \psi))$$

astrophysics

- Large uncertainties in the DM density profile in the inner few kpc.



# Dark Matter indirect detection

- Expected gamma-ray flux from DM annihilation:

$$\frac{d\Phi_\gamma}{dE} = \frac{\langle\sigma v\rangle}{8\pi m_\chi^2} \frac{dN_\gamma}{dE} \int_{\text{l.o.s.}} ds \overset{\text{astrophysics}}{\rho^2(r(s, \psi))}$$



- Large uncertainties in the DM density profile in the inner few kpc.

## Use cosmological simulations:

- DMO simulations predict NFW profile:  $r^{-\gamma}$ , where  $\gamma \approx 1$  in the inner few kpc.
- What is the DM density profile for **MW-like galaxies** in hydrodynamical simulations?

# Galactic centre GeV excess


- Unexplained excess of gamma rays in Fermi-LAT data from the centre of our Galaxy, above the known astrophysical background. Hooper & Goodenough '09, Vitale & Morselli '09, ...




Macias & Gordon, 1312.6671

- DM interpretation:**  
Best fit value for the inner slope:  $\gamma = 1.26 \pm 0.15$
- Other interpretations:** *unresolved millisecond pulsars, diffuse photons from cosmic rays, stellar source population in the Galactic bulge, ...*

# GeV excess DM interpretation



Calore et al., 1411.4647



Calore et al., 1409.0042

Generalized NFW:





$$\rho(r) = \rho_s \frac{r_s^3}{r^\gamma (r + r_s)^{3-\gamma}}$$

# GeV excess DM interpretation

- Test the DM density profile predicted by hydrodynamical simulations against the GeV excess data.
- **Additional selection criterion of MW-like galaxies:** substantial stellar disk component.





4 MW analogues:  
2 EAGLE + 2 APOSTLE

# Dark Matter density profiles



Calore, Bozorgnia et al., 1509.02164





# Dark Matter density profiles



Calore, Bozorgnia et al., 1509.02164



# Dark Matter density profiles



Calore, Bozorgnia et al., 1509.02164



# Dark Matter density profiles

- GeV excess data analyzed in the region:

$$2^\circ \leq |b| \leq 20^\circ \ \& \ |l| \leq 20^\circ$$

radial scale: 0.3 - 3 kpc

- **A very conservative approach:** power-law extrapolation with maximal asymptotic slope at the Power radius.




EAGLE HR (2 haloes):  $0.94 < \gamma_{\max} < 0.98$  at  $R_{P03} = 1.8$  kpc

APOSTLE IR (2 haloes):  $0.50 < \gamma_{\max} < 0.62$  at  $R_{P03} = 1.8$  kpc.

# Fitting the GeV excess

- Assuming 100% annihilation into b-quarks:





- Similar constraints on DM mass and annihilation cross section, but significantly worse fit.

(238 dof)

Profile	$\langle\sigma v\rangle [\times 10^{-26} \text{ cm}^3/\text{s}]$	$m_\chi$ [GeV]	$\chi^2$	$p$ -value
gNFW ( $\gamma=1.26$ )	$1.71 \pm 0.11$	$47.32 \pm 1.07$	223.9	0.73
EAGLE HR	$1.96 \pm 0.14$	$46.37 \pm 1.37$	246.3	0.34
APOSTLE IR	$1.76 \pm 0.16$	$45.36 \pm 2.96$	283.9	0.02

# Fitting the GeV excess

- Even under our very conservative assumption, DM density profiles of our MW-like galaxies do not reproduce the correct morphology of the GeV excess in the inner most regions.




Calore, Bozorgnia et al., 1509.02164

# Summary

- Need a precise determination of the DM distribution in the MW.  
→ *Identify MW analogues* in simulations by taking into account observational constraints on the MW.
- **Local DM density** agrees with local and global estimates.
- **DM density profiles** show flattening in the inner few kpc and contraction up to 10 kpc.
- **Halo integrals** match well those obtained from best fit Maxwellian velocity distributions.
- A **Maxwellian velocity distribution** with *peak speed* constrained by hydro simulations, and independent from the *local circular speed*, could be used for the analysis of direct detection data.
- DM density profiles of MW-like galaxies fail to reproduce the GeV excess.

# Backup Slides


# Selection criteria for MW analogues



- ▶  $M_*$  strongly correlated with  $v_c$  at 8 kpc, while the correlation of  $M_{200}$  with  $v_c$  is weaker.
- ▶  $M_*(R < 8 \text{ kpc}) = (0.5 - 0.9)M_*$ .
- ▶  $M_{\text{tot}}(R < 8 \text{ kpc}) = (0.01 - 0.1)M_{200}$ .
- ▶ Over the small halo mass range probed, little correlation between  $M_{\text{DM}}(R < 8 \text{ kpc})$  and  $M_{200}$ .

# Searching for dark disks

DM and stellar velocity distributions:



- ▶ Fit with a double Gaussian. Difference in the mean speed of second Gaussian between DM and stars is 35 km/s in the left, and 7 km/s in the right panel.
- ▶ Fraction of second Gaussian is 32% in the left panel and 43% in the right panel.



# Parameters of the simulations

Simulation	code	$N_{\text{DM}}$	$m_{\text{g}}$ [ $M_{\odot}$ ]	$m_{\text{DM}}$ [ $M_{\odot}$ ]	$\epsilon$ [pc]
Ling <i>et al.</i>	RAMSES	2662	–	$7.46 \times 10^5$	200
Eris	GASOLINE	81213	$2 \times 10^4$	$9.80 \times 10^4$	124
NIHAO	EFS-GASOLINE2	–	$3.16 \times 10^5$	$1.74 \times 10^6$	931
EAGLE (HR)	P-GADGET (ANARCHY)	1821–3201	$2.26 \times 10^5$	$1.21 \times 10^6$	350
APOSTLE (IR)	P-GADGET (ANARCHY)	2160, 3024	$1.3 \times 10^5$	$5.9 \times 10^5$	308
MaGICC	GASOLINE	4849, 6541	$2.2 \times 10^5$	$1.11 \times 10^6$	310
Sloane <i>et al.</i>	GASLOINE	5847–7460	$2.7 \times 10^4$	$1.5 \times 10^5$	174

## Properties of the selected MW analogues

Simulation	Count	$M_{\text{star}}$ [ $\times 10^{10} M_{\odot}$ ]	$M_{\text{halo}}$ [ $\times 10^{12} M_{\odot}$ ]	$\rho_{\chi}$ [ $\text{GeV}/\text{cm}^3$ ]	$v_{\text{peak}}$ [km/s]
Ling <i>et al.</i>	1	$\sim 8$	0.63	0.37–0.39	239
Eris	1	3.9	0.78	0.42	239
NIHAO	5	15.9	$\sim 1$	0.42	192–363
EAGLE (HR)	12	4.65–7.12	2.76–14.26	0.42–0.73	232–289
APOSTLE (IR)	2	4.48, 4.88	1.64–2.15	0.41–0.54	223–234
MaGICC	2	2.4–8.3	0.584, 1.5	0.346, 0.493	187, 273
Sloane <i>et al.</i>	4	2.24–4.56	0.68–0.91	0.3–0.4	185–204

# Morphology of simulated haloes

- ▶ Select simulated galaxies whose stellar kinematics show a disc component, rather than ellipticals or undergoing mergers.
- ▶ Characterize the morphology of each simulated galaxy by looking for evidence of coherent rotation.
- ▶ Use the distribution of angular momentum vectors of individual particles relative to the net angular momentum of the galaxy to discriminate between discs (coherent rotation) and spheroids (no coherent rotation).
- ▶ Derive the distribution of the stellar orbital circularity parameter,



$$\epsilon(r) = \frac{j_z}{j_c(r)}$$

A distribution peaked at  $\epsilon = 1 \Rightarrow$  disc


An almost symmetric distribution around  $\epsilon = 0 \Rightarrow$  spheroidal system

# Morphology of simulated haloes

- ▶ We retain a galaxy if the stellar fraction in the range  $\epsilon > 0.45$  is larger than 50%.
- ▶ With this criterion we can identify galaxies that have a dominant disc, and remove galaxies that show an almost symmetric distribution around  $\epsilon = 0$ .



# GeV excess DM interpretation



Channel	$\langle\sigma v\rangle$ ( $10^{-26} \text{ cm}^3 \text{ s}^{-1}$ )	$m_\chi$ (GeV)	$\chi^2_{\min}$	$p$ -value
$\bar{q}q$	$0.83^{+0.15}_{-0.13}$	$23.8^{+3.2}_{-2.6}$	26.7	0.22
$\bar{c}c$	$1.24^{+0.15}_{-0.15}$	$38.2^{+4.7}_{-3.9}$	23.6	0.37
$\bar{b}b$	$1.75^{+0.28}_{-0.26}$	$48.7^{+6.4}_{-5.2}$	23.9	0.35
$\bar{t}t$	$5.8^{+0.8}_{-0.8}$	$173.3^{+2.8}_{-0}$	43.9	0.003
$gg$	$2.16^{+0.35}_{-0.32}$	$57.5^{+7.5}_{-6.3}$	24.5	0.32
$W^+W^-$	$3.52^{+0.48}_{-0.48}$	$80.4^{+1.3}_{-0}$	36.7	0.026
$ZZ$	$4.12^{+0.55}_{-0.55}$	$91.2^{+1.53}_{-0}$	35.3	0.036
$hh$	$5.33^{+0.68}_{-0.68}$	$125.7^{+3.1}_{-0}$	29.5	0.13
$\tau^+\tau^-$	$0.337^{+0.047}_{-0.048}$	$9.96^{+1.05}_{-0.91}$	33.5	0.055
$[\mu^+\mu^-]$	$1.57^{+0.23}_{-0.23}$	$5.23^{+0.22}_{-0.27}$	43.9	0.0036] <del>ies</del>

Calore et al., 1411.4647